Justification Logic for Intuitionistic Modal Logic

Sonia Marin and Paaras Padhiar

School of Computer Science, University of Birmingham

September 2025

▶ In modal logic, $\Box A$ can mean A is provable or known.

- ▶ In modal logic, $\Box A$ can mean A is provable or known.
- ▶ Justification logic makes this knowledge explicit.

[Artemov]

- ▶ In modal logic, $\Box A$ can mean A is provable or known.
- ▶ Justification logic makes this knowledge explicit.

[Artemov]

- $ightharpoonup \Box A \leadsto t:A$, where t is a proof of A, or explicit knowledge of A.
- ► Read this as t justifies A.

▶ Proof terms are equipped with operations.

- ▶ Proof terms are equipped with operations.
- Gives an internal view of proof manipulations or tracing knowledge.

- Proof terms are equipped with operations.
- Gives an internal view of proof manipulations or tracing knowledge.
- ► E.g. Proof application internalising modus ponens:

$$k : \Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B) \rightsquigarrow jk : s:(A \rightarrow B) \rightarrow (t:A \rightarrow s \cdot t:B)$$

- Proof terms are equipped with operations.
- Gives an internal view of proof manipulations or tracing knowledge.
- ► E.g. Proof application internalising modus ponens:

$$k: \Box(A \to B) \to (\Box A \to \Box B) \leadsto jk: s:(A \to B) \to (t:A \to s \cdot t:B)$$

► Allow monotonicity of proofs

$$s_i:A \rightarrow s_1 + s_2:A$$

where $i \in \{1, 2\}$.

First justification logic, Logic of Proofs (LP) is corresponding logic for S4.
[Artemov]

$$S4 \vdash \Box A \iff \exists t \text{ such that } LP \vdash t:A^r$$

First justification logic, Logic of Proofs (LP) is corresponding logic for S4.
[Artemov]

$$S4 \vdash \Box A \iff \exists t \text{ such that } LP \vdash t:A^r$$

Provides a connection to PA:

$$\mathsf{LP} \vdash t : A \iff \mathsf{PA} \vdash \mathsf{Proof}(t, A^*)$$

First justification logic, Logic of Proofs (LP) is corresponding logic for S4.
[Artemov]

$$S4 \vdash \Box A \iff \exists t \text{ such that } LP \vdash t:A^r$$

Provides a connection to PA:

$$LP \vdash t:A \iff PA \vdash Proof(t, A^*)$$

- Other justification logics include:
 - ► Logics for S5 cube [Artemov et al., Brezhnev, Brünnler et al.]
 - Geach/Scott-Lemmon logics

[Fitting]

Gödel-Löb logic

[Shamkanov, Fitting]

► Non-normal modal logics

[Rohani and Studer]

▶ JL formally, extends the language of CPL with t:A.

- ▶ JL formally, extends the language of CPL with t:A.
- Proof terms $t := x \mid c \mid (t+t) \mid (t \cdot t)$.

- ▶ JL formally, extends the language of CPL with t:A.
- Proof terms $t := x \mid c \mid (t+t) \mid (t \cdot t)$.
 - x ranges over proof variables: x represents arbitrary proofs.

- ▶ JL formally, extends the language of CPL with t:A.
- ▶ Proof terms $t ::= x \mid c \mid (t+t) \mid (t \cdot t)$.
 - x ranges over proof variables: x represents arbitrary proofs.
 - c ranges over proof constants proof constants witness axioms.

- ▶ JL formally, extends the language of CPL with t:A.
- Proof terms $t := x \mid c \mid (t+t) \mid (t \cdot t)$.
 - x ranges over proof variables: x represents arbitrary proofs.
 - c ranges over proof constants proof constants witness axioms.
- Base axiomatisation extends CPL with

jk :
$$s:(A \rightarrow B) \rightarrow (t:A \rightarrow s \cdot t:B)$$

$$j+$$
 : $(s:A \lor t:A) \rightarrow s + t:A$

- ▶ JL formally, extends the language of CPL with t:A.
- Proof terms $t := x \mid c \mid (t+t) \mid (t \cdot t)$.
 - x ranges over proof variables: x represents arbitrary proofs.
 - c ranges over proof constants proof constants witness axioms.
- Base axiomatisation extends CPL with jk: $s:(A \rightarrow B) \rightarrow (t:A \rightarrow s \cdot t:B)$ j+: $(s:A \lor t:A) \rightarrow s + t:A$ $can \frac{A \text{ is an axiom instance}}{can \cdot (can \cdot Can \cdot Ca$
- Add other proof operators and axioms if required.

▶ Justification logic can internalise its own reasoning.

Justification logic can internalise its own reasoning.

Lemma (Lifting Lemma)

If $JL \vdash A_1 \rightarrow \ldots \rightarrow A_n \rightarrow B$, then there exists a proof term $t(x_1, \ldots, x_n)$ such that

$$\mathsf{LP} \vdash x_1 : A_1 \to \ldots \to x_n : A_n \to t(x_1, \ldots, x_n) : B$$

Justification logic can internalise its own reasoning.

Lemma (Lifting Lemma)

If $JL \vdash A_1 \rightarrow \ldots \rightarrow A_n \rightarrow B$, then there exists a proof term $t(x_1, \ldots, x_n)$ such that

$$\mathsf{LP} \vdash x_1 : A_1 \to \ldots \to x_n : A_n \to t(x_1, \ldots, x_n) : B$$

This mirrors the external modal view:

$$\mathsf{k} \; \frac{A_1, \ldots, A_n \Rightarrow B}{\Box A_1, \ldots, \Box A_n \Rightarrow \Box B}$$

► As a corollary, justification logic can emulate necessitation

$$nec \frac{\vdash A}{\vdash \Box A}$$

Corollary

If $JL \vdash A$, then there exists a proof term t such that

$$LP \vdash t:A$$

Both modal logic and justification logic share a similar structure.

- Both modal logic and justification logic share a similar structure.
- ► Terms behave externally like □:

- Both modal logic and justification logic share a similar structure.
- ► Terms behave externally like □:
 - ► Forgetting each term into a box
 - ▶ JL \vdash $A \Rightarrow$ ML \vdash A^f

- Both modal logic and justification logic share a similar structure.
- ► Terms behave externally like □:
 - ► Forgetting each term into a box
 - ightharpoonup JL \vdash $A \Rightarrow$ ML \vdash A^f
- ► For each □ we can find its corresponding proof term:

$$\mathsf{ML} \vdash \Box A \implies \exists t \text{ such that } \mathsf{JL} \vdash t : A^r$$

8 / 29

- Both modal logic and justification logic share a similar structure.
- ► Terms behave externally like □:
 - ► Forgetting each term into a box
 - ightharpoonup JL \vdash $A \Rightarrow$ ML \vdash A^f
- ► For each □ we can find its corresponding proof term:

$$\mathsf{ML} \vdash \Box A \implies \exists t \text{ such that } \mathsf{JL} \vdash t : A^r$$

.

ightharpoonup are realised into proof terms through a realisation function r.

- Both modal logic and justification logic share a similar structure.
- ► Terms behave externally like □:
 - ► Forgetting each term into a box
 - ightharpoonup JL $\vdash A \Rightarrow$ ML $\vdash A^{f}$
- ► For each □ we can find its corresponding proof term:

$$\mathsf{ML} \vdash \Box A \implies \exists t \text{ such that } \mathsf{JL} \vdash t : A^r$$

.

- □s are realised into proof terms through a realisation function r.
- Realisation can be done proof-theoretically or semantically.

[Artemov, Fitting]

Proof-theoretic realisation "re-writes" proofs of modal logic, into proofs of justification logic.

- ▶ Proof-theoretic realisation "re-writes" proofs of modal logic, into proofs of justification logic.
- ► First realisation method uses cut-free Gentzen-style sequent calculi. [Artemov, Brezhnev]

- ▶ Proof-theoretic realisation "re-writes" proofs of modal logic, into proofs of justification logic.
- First realisation method uses cut-free Gentzen-style sequent calculi. [Artemov, Brezhnev]
- Method was expanded to hypersequent and nested sequent calculi.

[Artemov et. al, Brünnler et. al]

- Proof-theoretic realisation "re-writes" proofs of modal logic, into proofs of justification logic.
- First realisation method uses cut-free Gentzen-style sequent calculi. [Artemov, Brezhnev]
- Method was expanded to hypersequent and nested sequent calculi.

[Artemov et. al, Brünnler et. al]

Briefly outline the method for constructing a realisation with nested sequents. [Goetschi and Kuznets]

Nested sequents

Nested sequents

- ► A nested sequent is a finite multiset of formulas and boxed sequents.
- $\blacktriangleright \ \Gamma ::= \varnothing \mid A, \Gamma \mid \Gamma, [\Gamma]$

Nested sequents

- ► A nested sequent is a finite multiset of formulas and boxed sequents.
- $ightharpoonup \Gamma ::= \varnothing \mid A, \Gamma \mid \Gamma, [\Gamma]$
- ▶ $fm(\emptyset) = \bot$, $fm(A, Γ) = A \lor fm(Γ)$, and $fm(Γ_1, [Γ_2]) = fm(Γ_1) \lor \Box fm(Γ_2)$

Nested sequents

- ► A nested sequent is a finite multiset of formulas and boxed sequents.
- $ightharpoonup \Gamma ::= \varnothing \mid A, \Gamma \mid \Gamma, [\Gamma]$
- ▶ $fm(\emptyset) = \bot$, $fm(A, \Gamma) = A \lor fm(\Gamma)$, and $fm(\Gamma_1, [\Gamma_2]) = fm(\Gamma_1) \lor \Box fm(\Gamma_2)$
- ► A context is a nested sequent with one or several holes { } which can take the place of a formula in the sequent.
- ▶ This lets us write $\Gamma_1\{\Gamma_2\}$ when we replace the hole in $\Gamma_1\{$ $\}$ by Γ_2

Nested sequents

- ► A nested sequent is a finite multiset of formulas and boxed sequents.
- $ightharpoonup \Gamma ::= \varnothing \mid A, \Gamma \mid \Gamma, [\Gamma]$
- ▶ $fm(\emptyset) = \bot$, $fm(A, \Gamma) = A \lor fm(\Gamma)$, and $fm(\Gamma_1, [\Gamma_2]) = fm(\Gamma_1) \lor \Box fm(\Gamma_2)$
- ► A context is a nested sequent with one or several holes { } which can take the place of a formula in the sequent.
- ▶ This lets us write $\Gamma_1\{\Gamma_2\}$ when we replace the hole in $\Gamma_1\{$ $\}$ by Γ_2
- ▶ E.g. $\Gamma\{\ \} = A, [B, [\{\ \}, [D]]]$. Then

$$\Gamma\{C\} = A, [B, [C, [D]]]$$

Nested system nK

$$\operatorname{id} \frac{\Gamma\{a, \overline{a}\}}{\Gamma\{a, \overline{a}\}} \qquad \wedge \frac{\Gamma\{A\} \quad \Gamma\{B\}}{\Gamma\{A \wedge B\}} \qquad \vee \frac{\Gamma\{A, B\}}{\Gamma\{A \vee B\}} \qquad \Box \frac{\Gamma\{[A]\}}{\Gamma\{\Box A\}}$$

$$\operatorname{k} \frac{\Gamma\{[A, \Delta]\}}{\Gamma\{\diamondsuit A, [\Delta]\}}$$

Nested system nK

$$\operatorname{id} \frac{\Gamma\{a,\bar{a}\}}{\Gamma\{a,\bar{a}\}} \qquad \wedge \frac{\Gamma\{A\} \quad \Gamma\{B\}}{\Gamma\{A \wedge B\}} \qquad \vee \frac{\Gamma\{A,B\}}{\Gamma\{A \vee B\}} \qquad \Box \frac{\Gamma\{[A]\}}{\Gamma\{\Box A\}}$$

$$\operatorname{k} \frac{\Gamma\{[A,\Delta]\}}{\Gamma\{\diamondsuit A,[\Delta]\}}$$

Theorem (Soundness and Completeness)

$$\mathsf{K} \vdash \mathsf{A} \iff \mathsf{n} \mathsf{K} \vdash \mathsf{A}$$

(Proved using a cut-elimination argument)

$$\operatorname{cut} \frac{\Gamma\{A\} \quad \Gamma\{\bar{A}\}}{\Gamma\{\varnothing\}}$$

Theorem

There exists a realisation function r such that

$$\mathsf{K} \vdash \mathsf{A} \Rightarrow \mathsf{J} \mathsf{K} \vdash \mathsf{A}^r$$

Theorem

There exists a realisation function r such that

$$\mathsf{K} \vdash \mathsf{A} \Rightarrow \mathsf{J} \mathsf{K} \vdash \mathsf{A}^r$$

Proof sketch:

► Take a cut-free nested sequent proof of *A*.

Theorem

There exists a realisation function r such that

$$\mathsf{K} \vdash \mathsf{A} \Rightarrow \mathsf{J} \mathsf{K} \vdash \mathsf{A}^r$$

Proof sketch:

- ► Take a cut-free nested sequent proof of A.
- Construct a realisation "top-down".

Theorem

There exists a realisation function r such that

$$\mathsf{K} \vdash \mathsf{A} \Rightarrow \mathsf{J} \mathsf{K} \vdash \mathsf{A}^r$$

Proof sketch:

- ► Take a cut-free nested sequent proof of A.
- ► Construct a realisation "top-down".
- Proceed by induction on the height of the proof of A.

id
$$\frac{}{\Gamma\{a,\bar{a}\}}$$

▶ For the base case, proof is an instance of id.

id
$$\frac{}{\Gamma\{a,\bar{a}\}}$$

▶ "Replicate" the soundness of this rule in justification logic.

$$\operatorname{id} \frac{}{\Gamma\{a,\bar{a}\}}$$

- ▶ "Replicate" the soundness of this rule in justification logic.
- ightharpoonup E.g. id $\frac{}{p,[a,\bar{a}]}$
- $\blacktriangleright \ \mathsf{fm}(p,[a,\bar{a}]) = p \vee \Box(a \vee \bar{a})$

$$\operatorname{id} \frac{}{\Gamma\{a,\bar{a}\}}$$

- "Replicate" the soundness of this rule in justification logic.
- $\blacktriangleright \text{ E.g. id } \frac{}{p,[a,\bar{a}]}$
- $\blacktriangleright \ \mathsf{fm}(p,[a,\bar{a}]) = p \vee \Box (a \vee \bar{a})$
- ► Construct a realisation through the following:

$$\operatorname{id} \frac{}{\Gamma\{a,\bar{a}\}}$$

- ▶ "Replicate" the soundness of this rule in justification logic.
- $\blacktriangleright \text{ E.g. id } \frac{}{p,[a,\bar{a}]}$
- $\blacktriangleright \ \mathsf{fm}(p,[a,\bar{a}]) = p \vee \Box(a \vee \bar{a})$
- ► Construct a realisation through the following:
- JK ⊢ a ∨ ā

$$\operatorname{id} \frac{}{\Gamma\{a,\bar{a}\}}$$

- ▶ "Replicate" the soundness of this rule in justification logic.
- $\blacktriangleright \text{ E.g. id } \frac{}{p,[a,\bar{a}]}$
- $\blacktriangleright \ \mathsf{fm}(p,[a,\bar{a}]) = p \vee \Box (a \vee \bar{a})$
- ► Construct a realisation through the following:
- JK ⊢ a ∨ ā
- ▶ Apply constructive necessitation JK $\vdash t:(a \lor \bar{a})$

$$\operatorname{id} \frac{}{\Gamma\{a,\bar{a}\}}$$

- "Replicate" the soundness of this rule in justification logic.
- $\blacktriangleright \text{ E.g. id } \frac{}{p,[a,\bar{a}]}$
- $\blacktriangleright \ \mathsf{fm}(p,[a,\bar{a}]) = p \vee \Box(a \vee \bar{a})$
- Construct a realisation through the following:
- JK ⊢ a ∨ ā
- ▶ Apply constructive necessitation JK $\vdash t:(a \lor \bar{a})$
- ▶ JK \vdash $p \lor t:(a \lor \bar{a})$

rule
$$\frac{\Gamma_1 \cdots \Gamma_n}{\Gamma}$$

For inductive case, consider each rule of the system:

rule
$$\frac{\Gamma_1 \cdots \Gamma_n}{\Gamma}$$

▶ Using the inductive hypothesis, we have some realisations r_1, \ldots, r_n with $\mathsf{JK} \vdash \Gamma_1^{r_1}, \ldots, \mathsf{JK} \vdash \Gamma_n^{r_n}$.

rule
$$\frac{\Gamma_1 \cdots \Gamma_n}{\Gamma}$$

- ▶ Using the inductive hypothesis, we have some realisations r_1, \ldots, r_n with $JK \vdash \Gamma_1^{r_1}, \ldots, JK \vdash \Gamma_n^{r_n}$.
- \triangleright Using these theorems, we construct a realisation r on Γ.

rule
$$\frac{\Gamma_1 \cdots \Gamma_n}{\Gamma}$$

- ▶ Using the inductive hypothesis, we have some realisations r_1, \ldots, r_n with $JK \vdash \Gamma_1^{r_1}, \ldots, JK \vdash \Gamma_n^{r_n}$.
- ightharpoonup Using these theorems, we construct a realisation r on Γ.
- "Replicate" the soundness of this rule in justification logic.

rule
$$\frac{\Gamma_1 \cdots \Gamma_n}{\Gamma}$$

- ▶ Using the inductive hypothesis, we have some realisations r_1, \ldots, r_n with $JK \vdash \Gamma_1^{r_1}, \ldots, JK \vdash \Gamma_n^{r_n}$.
- \triangleright Using these theorems, we construct a realisation r on Γ .
- "Replicate" the soundness of this rule in justification logic.
- Difficulty here is dealing with nesting where rules are applied within brackets.

Consider the following:

$$\wedge \frac{p,[q_1] \quad p,[q_2]}{p,[q_1 \wedge q_2]}$$

with realisations $(p,[q_1])^{r_1}=p\lor t_1{:}q_1$ and $(p,[q_2])^{r_2}=p\lor t_2{:}q_2$

Consider the following:

$$\wedge \frac{p,[q_1] \quad p,[q_2]}{p,[q_1 \wedge q_2]}$$

with realisations $(p,[q_1])^{r_1} = p \vee t_1:q_1$ and $(p,[q_2])^{r_2} = p \vee t_2:q_2$

Construct the new realisation *r* with the following:

Consider the following:

$$\wedge \frac{p,[q_1] \quad p,[q_2]}{p,[q_1 \wedge q_2]}$$

with realisations $(p,[q_1])^{r_1}=p\lor t_1:q_1$ and $(p,[q_2])^{r_2}=p\lor t_2:q_2$

- Construct the new realisation r with the following:
 - $\blacktriangleright \;\; \mathsf{JK} \vdash q_1 \to q_2 \to (q_1 \land q_2)$

Consider the following:

$$\wedge \, \frac{p,[q_1] \quad p,[q_2]}{p,[q_1 \wedge q_2]}$$

with realisations $(p,[q_1])^{r_1}=p\lor t_1:q_1$ and $(p,[q_2])^{r_2}=p\lor t_2:q_2$

- Construct the new realisation r with the following:
 - ightharpoonup JK $\vdash q_1 \rightarrow q_2 \rightarrow (q_1 \land q_2)$
 - ▶ Lifting Lemma: JK $\vdash t_1:q_1 \rightarrow t_2:q_2 \rightarrow t(t_1,t_2):(q_1 \land q_2)$

Consider the following:

$$\wedge \frac{p,[q_1] \quad p,[q_2]}{p,[q_1 \wedge q_2]}$$

with realisations $(p,[q_1])^{r_1}=p\lor t_1:q_1$ and $(p,[q_2])^{r_2}=p\lor t_2:q_2$

- Construct the new realisation r with the following:
 - ightharpoonup JK $\vdash q_1 \rightarrow q_2 \rightarrow (q_1 \land q_2)$
 - ▶ Lifting Lemma: JK $\vdash t_1:q_1 \rightarrow t_2:q_2 \rightarrow t(t_1,t_2):(q_1 \land q_2)$
 - $\blacktriangleright \ \mathsf{JK} \vdash (p \lor t_1 \mathpunct{:} q_1) \mathbin{\rightarrow} (p \lor t_2 \mathpunct{:} q_2) \mathbin{\rightarrow} (p \lor t(t_1, t_2) \mathpunct{:} (q_1 \land q_2))$

Rules give a manageable way constructing realisations top-down.

- Rules give a manageable way constructing realisations top-down.
- ► Modus ponens and cut rules in modal logic don't always translate to justification logic.

- Rules give a manageable way constructing realisations top-down.
- Modus ponens and cut rules in modal logic don't always translate to justification logic.
- ► E.g.

$$\operatorname{mp} \frac{\Box A \to \Box A \quad (\Box A \to \Box A) \to B}{B}$$

- Rules give a manageable way constructing realisations top-down.
- Modus ponens and cut rules in modal logic don't always translate to justification logic.
- ► E.g.

$$\operatorname{mp} \frac{\Box A \to \Box A \quad (\Box A \to \Box A) \to B}{B}$$

$$\operatorname{mp} \frac{x:A \to t \cdot x:A \quad (x:A \to x:A) \to B}{?}$$

where *t* is a proof of $A \rightarrow A$.

► Focus is on intuitionistic justification logics.

- Focus is on intuitionistic justification logics.
- Past work includes:
 - Explicit counterpart to iS4.

[Artemov]

- Focus is on intuitionistic justification logics.
- Past work includes:
 - Explicit counterpart to iS4.

[Artemov]

Connections to Heyting arithmetic (HA)

[Artemov and lemhoff, Dashkov]

- ► Focus is on intuitionistic justification logics.
- Past work includes:
 - Explicit counterpart to iS4. [Artemov]
 - ► Connections to Heyting arithmetic (HA)

[Artemov and lemhoff, Dashkov]

Curry-Howard correspondence

[Steren and Bonelli]

- ► Focus is on intuitionistic justification logics.
- Past work includes:
 - Explicit counterpart to iS4. [Artemov]
 - ► Connections to Heyting arithmetic (HA)

[Artemov and lemhoff, Dashkov]

Curry-Howard correspondence

[Steren and Bonelli]

Intuitionistic diamonds

► What about ♦?

- ▶ What about ♦?
- ▶ Intuitionistic \Box and \Diamond are not De Morgan dual.

$$\Box A \not\leftrightarrow \neg \Diamond \neg A$$

- ► What about ♦?
- ▶ Intuitionistic □ and ⋄ are not De Morgan dual.

$$\Box A \not\leftrightarrow \neg \Diamond \neg A$$

▶ Language \mathcal{L}_{\square} :

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \to A) \mid \Box A \mid \Diamond A$$

- ► What about ♦?
- ▶ Intuitionistic □ and ⋄ are not De Morgan dual.

$$\Box A \not\leftrightarrow \neg \Diamond \neg A$$

▶ Language \mathcal{L}_{\square} :

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \to A) \mid \Box A \mid \Diamond A$$

► Smallest intuitionistic modal logic with ♦ is constructive modal logic CK. [Bellin et al.]

$$k_1: \Box(A \to B) \to (\Box A \to \Box B)$$

$$k_2: \Box(A \to B) \to (\Diamond A \to \Diamond B)$$

- ► What about ♦?
- ▶ Intuitionistic □ and ⋄ are not De Morgan dual.

$$\Box A \not\leftrightarrow \neg \Diamond \neg A$$

▶ Language \mathcal{L}_{\square} :

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \to A) \mid \Box A \mid \Diamond A$$

Smallest intuitionistic modal logic with ♦ is constructive modal logic CK. [Bellin et al.]

$$\mathsf{k}_1:\Box(A\to B)\to(\Box A\to\Box B)$$

$$k_2: \Box(A \to B) \to (\Diamond A \to \Diamond B)$$

CK (and some extensions) have Gentzen-style proof theory

[Bierman and de Paiva]

► Idea: Make diamond explicit with satisfiers.

$$\Diamond A \leadsto \mu : A$$

 \blacktriangleright μ :A read as μ satisfies A, or μ is a model of A.

[Kuznets, Marin, Straßburger]

► Language *L*_J:

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \to A) \mid t:A \mid \mu:A$$

Idea: Make diamond explicit with satisfiers.

$$\Diamond A \rightsquigarrow \mu : A$$

 \blacktriangleright μ : A read as μ satisfies A, or μ is a model of A.

[Kuznets, Marin, Straßburger]

► Language L_J:

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \to A) \mid t:A \mid \mu:A$$

► Satisfier terms $\mu ::= \alpha \mid (\mu \sqcup \mu) \mid (t \star \mu)$

Idea: Make diamond explicit with satisfiers.

$$\Diamond A \leadsto \mu : A$$

 \blacktriangleright μ :A read as μ satisfies A, or μ is a model of A.

[Kuznets, Marin, Straßburger]

► Language L_J:

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \to A) \mid t : A \mid \mu : A$$

- ► Satisfier terms $\mu ::= \alpha \mid (\mu \sqcup \mu) \mid (t \star \mu)$
- Proof terms encapsulates global reasoning.

Idea: Make diamond explicit with satisfiers.

$$\Diamond A \leadsto \mu : A$$

 \blacktriangleright μ :A read as μ satisfies A, or μ is a model of A.

[Kuznets, Marin, Straßburger]

► Language L_J:

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \to A) \mid t : A \mid \mu : A$$

- ► Satisfier terms $\mu ::= \alpha \mid (\mu \sqcup \mu) \mid (t \star \mu)$
- Proof terms encapsulates global reasoning.
- Satisfier terms encapsulates local reasoning, or reasoning about consistency.

Justification counterpart: [Kuznets, Marin and Straßburger] $jk_1: s:(A \rightarrow B) \rightarrow (t:A \rightarrow s \cdot t:B)$

Justification counterpart: [Kuznets, Marin and Straßburger] jk₁ : $s:(A \rightarrow B) \rightarrow (t:A \rightarrow s \cdot t:B)$ jk₂ : $s:(A \rightarrow B) \rightarrow (\mu:A \rightarrow s \star \mu:B)$

Method of realisation is established similarly.

 Objective: justification counterpart of intuitionistic variant of modal logic IK [Fischer Servi, Plotkin and Stirling]

- Objective: justification counterpart of intuitionistic variant of modal logic IK [Fischer Servi, Plotkin and Stirling]
- ► Has following axiomatisation by extending IPL with:

 k_1 : $\Box(A \to B) \to (\Box A \to \Box B)$ k_2 : $\Box(A \to B) \to (\Diamond A \to \Diamond B)$

- Objective: justification counterpart of intuitionistic variant of modal logic IK [Fischer Servi, Plotkin and Stirling]
- Has following axiomatisation by extending IPL with:

```
\begin{array}{lll} \mathsf{k}_1 & : & \Box(A \to B) \to (\Box A \to \Box B) \\ \mathsf{k}_2 & : & \Box(A \to B) \to (\Diamond A \to \Diamond B) \\ \mathsf{k}_3 & : & \Diamond(A \lor B) \to (\Diamond A \lor \Diamond B) \end{array}
```

 $k_4 : (\Diamond A \rightarrow \Box B) \rightarrow \Box (A \rightarrow B)$

 k_5 : $\diamondsuit \bot \to \bot$

- Objective: justification counterpart of intuitionistic variant of modal logic IK [Fischer Servi, Plotkin and Stirling]
- ► Has following axiomatisation by extending IPL with:

$$\begin{array}{rcl} \mathsf{k}_1 & : & \Box(A \to B) \to (\Box A \to \Box B) \\ \mathsf{k}_2 & : & \Box(A \to B) \to (\diamondsuit A \to \diamondsuit B) \\ \mathsf{k}_3 & : & \diamondsuit(A \lor B) \to (\diamondsuit A \lor \diamondsuit B) \\ \mathsf{k}_4 & : & (\diamondsuit A \to \Box B) \to \Box(A \to B) \\ \mathsf{k}_5 & : & \diamondsuit \bot \to \bot \\ & & \mathsf{nec} \, \frac{\vdash A}{\vdash \Box A} \end{array}$$

▶ Language \mathcal{L}_J :

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \rightarrow A) \mid t:A \mid \mu:A$$

▶ Language \mathcal{L}_J :

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \rightarrow A) \mid t:A \mid \mu:A$$

▶ Proof terms $t ::= x \mid c \mid (t+t) \mid (t \cdot t) \mid (\mu \triangleright t)$.

► Language L_J:

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \rightarrow A) \mid t:A \mid \mu:A$$

- ▶ Proof terms $t ::= x \mid c \mid (t+t) \mid (t \cdot t) \mid (\mu \triangleright t)$.
- ► Satisfier terms $\mu ::= \alpha \mid (\mu \sqcup \mu) \mid (t \star \mu)$

► Language L_J:

$$A ::= \bot \mid p \mid (A \land A) \mid (A \lor A) \mid (A \rightarrow A) \mid t:A \mid \mu:A$$

- ▶ Proof terms $t ::= x \mid c \mid (t+t) \mid (t \cdot t) \mid (\mu \triangleright t)$.
- ▶ Satisfier terms $\mu ::= \alpha \mid (\mu \sqcup \mu) \mid (t \star \mu)$

$$\begin{aligned} \mathsf{jk_1} & : & s{:}(A \to B) \to (t{:}A \to s \cdot t{:}B) \\ \mathsf{jk_2} & : & s{:}(A \to B) \to (\mu{:}A \to s \star \mu{:}B) \end{aligned}$$

$$\begin{aligned} \mathsf{jk}_1 &: & s{:}(A \to B) \to (t{:}A \to s \cdot t{:}B) \\ \mathsf{jk}_2 &: & s{:}(A \to B) \to (\mu{:}A \to s \star \mu{:}B) \\ \mathsf{jk}_3 &: & \mu{:}(A \lor B) \to (\mu{:}A \lor \mu{:}B) \end{aligned}$$

$$\begin{aligned} \mathsf{jk_1} &: & \mathsf{s} \colon (A \to B) \to (t \colon A \to \mathsf{s} \cdot t \colon B) \\ \mathsf{jk_2} &: & \mathsf{s} \colon (A \to B) \to (\mu \colon A \to \mathsf{s} \star \mu \colon B) \\ \mathsf{jk_3} &: & \mu \colon (A \lor B) \to (\mu \colon A \lor \mu \colon B) \\ \mathsf{jk_4} &: & (\mu \colon A \to t \colon B) \to \mu \rhd t \colon (A \to B) \end{aligned}$$

$$\begin{array}{lll} \mathbf{j} \mathbf{k}_1 & : & s \colon (A \to B) \to (t \colon A \to s \cdot t \colon B) \\ \mathbf{j} \mathbf{k}_2 & : & s \colon (A \to B) \to (\mu \colon A \to s \star \mu \colon B) \\ \mathbf{j} \mathbf{k}_3 & : & \mu \colon (A \lor B) \to (\mu \colon A \lor \mu \colon B) \\ \mathbf{j} \mathbf{k}_4 & : & (\mu \colon A \to t \colon B) \to \mu \rhd t \colon (A \to B) \\ \mathbf{j} \mathbf{k}_5 & : & \mu \colon \bot \to \bot \end{array}$$

$$\begin{array}{rcl} \mathrm{jk_1} & : & s:(A \to B) \to (t:A \to s \cdot t:B) \\ \mathrm{jk_2} & : & s:(A \to B) \to (\mu:A \to s \star \mu:B) \\ \mathrm{jk_3} & : & \mu:(A \lor B) \to (\mu:A \lor \mu:B) \\ \mathrm{jk_4} & : & (\mu:A \to t:B) \to \mu \rhd t:(A \to B) \\ \mathrm{jk_5} & : & \mu:\bot \to \bot \\ \mathrm{j+} & : & (s:A \lor t:A) \to s + t:A \\ \mathrm{j} \sqcup & : & (\mu:A \lor \nu:A) \to \mu \sqcup \nu:A \\ & & \underbrace{A \text{ is an axiom instance}}_{G_0: \ldots:G_1:A} \end{array}$$

- ightharpoonup Each proof term replaced with a \Box .
- ► Each satisfier term replaced with a ⋄.

- ightharpoonup Each proof term replaced with a \Box .
- ► Each satisfier term replaced with a ♦.
- ► Achieved through the following:

Definition (Forgetful projection)

The forgetful projection is a map

$$(\cdot)^{\mathsf{f}}:\mathcal{L}_{\mathsf{J}}
ightarrow\mathcal{L}_{\square}$$

inductively defined as follows:

$$\begin{array}{ccc} \bot^{\mathsf{f}} & := \bot & & (t:A)^{\mathsf{f}} & := \Box A^{\mathsf{f}} \\ p^{\mathsf{f}} & := p & & (\mu:A)^{\mathsf{f}} & := \diamondsuit A^{\mathsf{f}} \\ (A*B)^{\mathsf{f}} & := (A^{\mathsf{f}}*B^{\mathsf{f}}) \text{ where } * \in \{\land, \lor, \to\} \end{array}$$

Result is formally stated as follows:

Theorem

 $\mathsf{JIK} \vdash A \Rightarrow \mathsf{IK} \vdash A^{\mathsf{f}}$.

Proof.

Apply forgetful projection on axioms of JIK to see we get theorems of IK.

For example:

$$\mathsf{jk}_4: ((\mu:A \to t:B) \to \mu \rhd t:(A \to B))^\mathsf{f} = (\Diamond A \to \Box B) \to \Box (A \to B)$$

25 / 29

Idea:

► Systematically replace each □ with a proof term, and each ⋄ with a satisfier term.

Idea:

- ► Systematically replace each □ with a proof term, and each ⋄ with a satisfier term.
- ► The objective is to produce a realisation function:

Idea:

- Systematically replace each □ with a proof term, and each ◇ with a satisfier term.
- ► The objective is to produce a realisation function:

Definition (Realisation)

A realisation is a map $(\cdot)^r: \mathcal{L}_{\square} \to \mathcal{L}_{\mathsf{J}}$ such that $(A^r)^{\mathsf{f}} = A$ for each $A \in \mathcal{L}_{\square}$.

Correspondence is formally stated as:

Theorem (Marin and P.)

There exists a realisation function r such that

$$\mathsf{IK} \vdash \mathsf{A} \Rightarrow \mathsf{JIK} \vdash \mathsf{A}^r$$

Correspondence is formally stated as:

Theorem (Marin and P.)

There exists a realisation function r such that

$$\mathsf{IK} \vdash A \Rightarrow \mathsf{JIK} \vdash A^r$$

▶ Only cut-free system available is a nested sequent system.

[Straßburger]

Correspondence is formally stated as:

Theorem (Marin and P.)

There exists a realisation function r such that

$$\mathsf{IK} \vdash A \Rightarrow \mathsf{JIK} \vdash A^r$$

Only cut-free system available is a nested sequent system.

[Straßburger]

A two-sided system with both structural boxes and diamonds.

Correspondence is formally stated as:

Theorem (Marin and P.)

There exists a realisation function r such that

$$\mathsf{IK} \vdash A \Rightarrow \mathsf{JIK} \vdash A^r$$

Only cut-free system available is a nested sequent system.

[Straßburger]

- ► A two-sided system with both structural boxes and diamonds.
- Adapt the method in the classical case.

Provides a treatment to make ◊ explicit for most intuitionistic modal logics.


- Provides a treatment to make ◊ explicit for most intuitionistic modal logics.
- Method for realisation can be applied to other nested sequent systems for intuitionistic modal logics.

[Arisaka et al., Kuznets and Straßburger]

Future work:


Future work:

Exploring connections to arithmetic.



Future work:


Exploring connections to arithmetic.



▶ Understanding ♦ and satisfiers in classical logic.

Future work:

Exploring connections to arithmetic.



▶ Understanding ♦ and satisfiers in classical logic.

- Semantics for JIK.
- Proof theory of JIK